Student Medical Exemption Certificate for Required Immunizations

Name of Primary Care Provider granting exemption:		
Please check one (practitioner granting exemption must be licensed as one of the following):		
🗌 Physician (MD or DO) 🗌 Physician Assistant	□ APRN	
CT License number:		
<u>NPI:</u>		
Phone number:	Email:	

Directions:

Part 1. Please complete the demographics section on the patient/student.

Part 2. Please mark the contraindications/precautions that apply to this patient/student (indicate all that apply).

Part 3. If no contraindications or precautions apply in part 2, write a brief explanation of the reason the patient/student requires the exemption.

Part 4. Sign the Statement of Clinical Opinion and date the form.

Attach a copy of the patient/student's most current immunization record.

Part 1. Patient/Student Information:

First name <u>(in full)</u>	Middle initial	Last name	
Date of Birth			
Mailing Address	City	·	
State			
Parent/Guardian: First Name		Last name	
Primary phone number			
School name			
School address			
City			
State	Zip		
Current or Grade student is entering			

Part 2. Please mark the vaccine(s), exemption duration, and all contraindications/precautions that apply to this patient/student for each vaccine.

Medical contraindications and precautions for immunizations are based upon the Advisory Committee on Immunization Practices (ACIP) <u>Comprehensive General Recommendations and Guidelines</u>, published by the Centers for Disease Control and Prevention.

A **contraindication** is a condition in a recipient that increases the risk for a serious vaccine adverse event (VAE) or compromises the ability of the vaccine to produce immunity.

A **precaution** is a condition in a recipient that might increase the risk for a serious VAE or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations are deferred when a precaution is self-limiting, but can be administered if the precaution condition improves.

Vaccine	Exemption Duration	ACIP Contraindications and Precautions (Check all that apply)
□ Diphtheria- Tetanus-and acellular	Temporary	Contraindications
	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	
Pertussis (DTaP)	mm/ yyyy	Encephalopathy within seven days after receipt of previous dose
(DTAF)	Permanent	of DTP or DTaP
		Precautions
		Progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy: defer DTaP until neurologic status clarified and stabilized
		 GBS <6 weeks after previous dose of tetanus-toxoid–containing vaccine
		Fever greater than 40.5°C (104.9°F) <48 hours after vaccination of previous dose of DTP or DTaP
		 History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid–containing or tetanus-toxoid– containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid–containing vaccine
		Moderate or acute illness with or without fever
Hepatitis A	Temporary	Contraindications
through: / mm/ yyyy □ Permanent	□ Severe allergic reaction (e.g., anaphylaxis) after a previous dose	
	/	or to a vaccine component
	_	Precautions
	□ Permanent	Moderate or severe acute illness with or without fever

CDC Recognized Contraindications and Precautions

	1	
□ Hepatitis B □ Tempo through: /	Temporary	Contraindications
	/	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	Hypersensitivity to yeast
	Permanent	Precautions
		Moderate or severe acute illness with or without fever
□Haemophilus		Contraindications
<i>influenza</i> e type b (HiB)		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
		□ Age <6 weeks
		Precautions
		Moderate or severe acute illness with or without fever
□ Inactivated	□ Temporary through: / mm/ yyyy □ Permanent	Contraindications
Influenza Virus ti (IIV)		Severe allergic reaction (e.g., anaphylaxis) after previous dose of influenza vaccine or to vaccine component
		Precautions
		\Box GBS <6 weeks after a previous dose of influenza vaccine
		Moderate or severe acute illness with or without fever
		Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a health care provider who is able to recognize and manage severe allergic conditions).
Polio Vaccine the (IPV) m	□ Temporary through: / mm/ yyyy □ Permanent	Contraindications
		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
		Precautions
		Pregnancy
		Moderate or acute illness with or without fever

Attenuated throug Influenza Virus (LAIV)	Temporary through:/	Contraindications
		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm∕ yyyy □ Permanent	 Concomitant use of aspirin or aspirin-containing medication in children and adolescents
		 LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days.(e)
		Pregnancy
		Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months.
		 Persons with active cerebrospinal fluid/oropharyngeal communications/leaks.
		Close contacts and caregivers of severely immunosuppressed persons who require a protected environment.
		 Persons with cochlear implants (due to the potential for CSF leak, which might exist for some period of time after implantation. Providers might consider consultation with a specialist concerning risk of persistent CSF leak if an age-appropriate inactivated or recombinant vaccine cannot be used).
		Altered Immunocompetence
		□ Anatomic or functional asplenia (e.g. sickle cell disease)
		Precautions
		□ GBS <6 weeks after a previous dose of influenza vaccine
		Asthma in persons aged 5 years old or older
		 Medical conditions which might predispose to higher risk of complications attributable to influenza(d)
		Moderate or severe acute illness with or without fever
🗆 Meningo-	□ Temporary through: / mm/ yyyy	Contraindications
coccal conjugate		Severe allergic reaction (e.g., anaphylaxis) after a previous dose
vaccines (MenACWY)		or to a vaccine component, including yeast Precautions
	Permanent	Moderate or severe acute illness with or without fever

	Temporary through:/	Contraindications
		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	Pregnancy
□ P	Permanent	Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised)
		□ Family history of altered immunocompetence (i)
		Precautions
		□ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product)
		□ History of thrombocytopenia or thrombocytopenic purpura
		 Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing (k)
		Moderate or severe acute illness with or without fever
🗆 Pneumo-	Temporary	Contraindications
	through: / mm/ yyyy Permanent	Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid–containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid– containing vaccine), including yeast
		Precautions
		Moderate or acute illness with or without fever
🗆 Tdap	Temporary	Contraindications
-	through: /	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or Tdap
		Precautions
		□ GBS <6 weeks after a previous dose of tetanus-toxoid–containing vaccine
		Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized
		□ History of Arthus-type hypersensitivity reactions after a previous

		 dose of diphtheria-toxoid—containing or tetanus-toxoid—containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine Moderate or severe acute illness with or without fever
□ Varicella	 □ Temporary through: / mm/ yyyy □ Permanent 	 Contraindications Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) (g) Pregnancy Family history of altered immunocompetence (j) Precautions Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product)
		Moderate or acute illness with or without fever

Part 3. Other Allergic Reactions/ Other Type of Medical Condition

Complete this section if claiming a medical exemption for a vaccine based on a condition that does meet any of the ACIP criteria for a contraindication or precaution listed in part 2.

Vaccine(s), list all that apply: _____

For each vaccine listed above, select the allergic or other reaction for which medical exemption is being submitted. Please check off any of the following that apply:

- □ This patient has an autoimmune disorder
- □ This patient has a family history of an autoimmune disorder
- □ This patient has a family history of a reaction to a vaccination
- □ This patient has a genetic predisposition to a reaction to a vaccination as determined through genetic testing
- □ This patient has a previous documented reaction that is correlated to a vaccination
- Other condition/reaction not listed above (must specify): ______

Please provide an explanation of the reaction/condition listed above:

Part 4. Statement of Clinical Opinion

In accord with the legal requirements of Public Act 21-6, the vaccine(s) indicated above is/are in my clinical opinion medically contraindicated for this patient/student due to the physical condition as explained above.

Clinician's Signature_____

Date _____

A person may be placed into quarantine or isolation when there are "reasonable grounds to believe [a person] to be infected with, or exposed to, a communicable disease or to be contaminated or exposed to contamination or at reasonable risk of having a communicable disease or being contaminated or passing such communicable disease or contamination to other persons if the commissioner determines that such individual or individuals pose a significant threat to the public health and that quarantine or isolation is necessary and the least restrictive alternative to protect or preserve the public health." Conn. Gen. Stat. § 19a-131b(a).